



Subsurface Scattering & Complex Material Properties

Last Time?

- What is a Pixel?
- Aliasing
- Fourier Analysis
- Sampling & Reconstruction
- Mip maps



Final Projects

- Teams of 2 highly encouraged
- Individuals or teams > 2 must talk to me first
- Continue to discuss on LMS
- Proposals due next week (Thursday 4/8)
 - Proposed project summary
 - At least 3 related papers (read & summarized)
 - Description of series of test cases
 - Timeline & initial task assignment

Today

- **Measuring BRDFs**
- 3D Digitizing & Scattering
- Complex Material Properties
- Importance of Participating Media
- BSSRDFs
- Other Complex Materials

BRDFs in the Movie Industry


- Agent Smith's clothes are CG, with measured BRDF



<http://www.virtualcinematography.org/publications/acrobat/BRDF-s2003.pdf>


How Do We Obtain BRDFs?

- Gonioreflectometer
 - 4 degrees of freedom




Source: Greg Ward

BRDFs in the Movie Industry



- <http://www.virtualcinematography.org/publications/acrobat/BRDF-s2003.pdf>

BRDFs in the Movie Industry




Not just a BRDF...




From presentation by
J.P. Lewis and George Borshukov

Materials – BRDF & BTDF



M. Andersen, "Innovative bi-directional video-goniophotometer for advanced fenestration systems", 2004.

Measuring Materials



M. Andersen, "Innovative bi-directional video-goniophotometer for advanced fenestration systems", 2004.


Today

- Measuring BRDFs
- **3D Digitizing & Scattering**
- Complex Material Properties
- Importance of Participating Media
- BSSRDFs
- Other Complex Materials

3D Digitizing



Cyberware



*The Digital Michelangelo Project:
3D Scanning of Large Statues,
Levoy et al., SIGGRAPH 2000*

Scattering & Scanning




Figure 1: Diffusion in a sample of Carrara Statuario marble.

*An Assessment of Laser Range Measurement
of Marble Surfaces, Godin et al, 2001.*

Questions?

Reading for Today:

- “Radiance Caching for Participating Media”, Jarosz, Donner, Zwicker, & Jensen, 2008.




Today

- Measuring BRDFs
- 3D Digitizing & Scattering
- **Complex Material Properties**
- **Importance of Participating Media**
- BSSRDFs
- Other Complex Materials

Anisotropic BRDFs

- Surfaces with strongly oriented microgeometry
- Examples:
 - brushed metals, hair, fur, cloth, velvet



Source: Westin et al 92

What makes a Rainbow?

- Refraction is wavelength-dependent
 - Refraction increases as the wavelength of light decreases
 - violet and blue experience more bending than orange and red
- Usually ignored in graphics
- Rainbow is caused by refraction + internal reflection + refraction






Pink Floyd, *The Dark Side of the Moon*



From "Color and Light in Nature" by Lynch and Livingstone

Amount of Reflection

- Traditional ray tracing (hack)
 - Constant `reflectionColor`
- More realistic:
 - Fresnel reflection term (more reflection at grazing angle)
 - Schlick's approximation: $R(\theta) = R_0 + (1 - R_0)(1 - \cos \theta)^5$



Dusty Surfaces & Retro-Reflection

- Viewed perpendicular to the surface, there is little scattering off dust
- At grazing angles, there is increased scattering with the dust making the surface appear brighter
- Similarly, the earth viewed from space appears brighter near the edges, because of increased scattering of the atmosphere.






Figure 5: Showing retroreflection from a very rough surface (left). Only areas with normals close to the light direction are well lit, so there is a strong retroreflective peak. On the right, we see a corner reflector (the inside corner of 3 planes is the 3D analog) which produces the same effect.

Light Rays in a Dusty Room



Annie Ding, MIT 6.837 Final Project December, 2004

Ray Tracing Participating Media



Participating Media





Image by Henrik Wann Jensen

Today

- Measuring BRDFs
- 3D Digitizing & Scattering
- Complex Material Properties
- Importance of Participating Media
- **BSSRDFs**
- Other Complex Materials

BRDF vs. BSSRDF



Images from "A Practical Model for Subsurface Light Transport"
Jensen, Marschner, Levoy, & Hanrahan SIGGRAPH 2001

Sampling a BSSRDF




Figure 7: (a) Sampling a BRDF (traditional sampling), (b) sampling a BSSRDF (the sample points are distributed both over the surface as well as the light).

Images from "A Practical Model for Subsurface Light Transport"
Jensen, Marschner, Levoy, & Hanrahan SIGGRAPH 2001

"A Practical Model for Subsurface Light Transport", Jensen, Marschner, Levoy, & Hanrahan, SIGGRAPH 2001




Subsurface Scattering Variables

Name	Symbol	Units	Description
Scattering Coeff.	σ_s	$(\text{length})^{-1}$	Probability of scattering per unit length
Absorption Coeff.	σ_a	$(\text{length})^{-1}$	Probability of absorption per unit length
Phase Function	$p(x, \vec{\omega}', \vec{\omega})$		Angular distribution of scattering
Extinction Coeff.	σ_t	$(\text{length})^{-1}$	$\sigma_a + \sigma_s$
(Scattering) Albedo	A		$\frac{\sigma_s}{\sigma_t}$
Optical Depth	$\tau(0, d)$		$\int_0^d \sigma_t dx$
Transmittance	$t(0, d)$		$e^{-\tau(0, d)}$

- Albedo: first approximation of BRDF, % of light reflected off the surface
 - When the albedo = 1, no absorption occurs and light is only transmitted or scattered. This is an ok approximation for snow or clouds.

BSSRDF Measurement



Images from "A Practical Model for Subsurface Light Transport"
Jensen, Marschner, Levoy, & Hanrahan SIGGRAPH 2001

Single Scattering




Figure 4: Single scattering occurs only when the refracted incoming and outgoing rays intersect, and is computed as an integral over path length s along the refracted outgoing ray.

Images from "A Practical Model for Subsurface Light Transport"
Jensen, Marschner, Levoy, & Hanrahan SIGGRAPH 2001

Dipole Approx. for Diffuse Scattering





Figure 3: An incoming ray is transformed into a dipole source for the diffusion approximation.

Images from "A Practical Model for Subsurface Light Transport"
Jensen, Marschner, Levoy, & Hanrahan SIGGRAPH 2001

Today

- Measuring BRDFs
- 3D Digitizing & Scattering
- Complex Material Properties
- Importance of Participating Media
- BSSRDFs
- **Other Complex Materials**




Measuring BSSRDF by Dilution

"Acquiring Scattering Properties of Participating Media by Dilution"
Narasimhan et al. SIGGRAPH 2006



Measuring Hair



"Light Scattering from Human Hair Fibers"
Marschner et al., SIGGRAPH 2003

Rendering Hair



Figure 12: A comparison of Kajiy and Kay's model (left) under a single point source, our proposed model (center) with the same lighting, and the hair from the photograph in Figure 11 (removed from context to simplify the comparison). The Kajiy model's diffuse term results in a flat appearance, while the secondary highlight in our model correctly captures the colored shading of the real hair.

"Light Scattering from Human Hair Fibers"
Marschner et al., SIGGRAPH 2003

Reading for Friday 4/1:

- "Ray Tracing on Programmable Graphics Hardware Purcell", Buck, Mark, & Hanrahan SIGGRAPH 2002

