



Announcements

- <u>Final exam schedule</u> has been posted by the Registrar. We are on M 4/28, 11:30am. (Room(s) TBD.)
- Exam 2 is this Wednesday 2/26. No recitations that day.
 - Last names A-K DCC 318; last names L-Z DCC 324.
 - One-page (double-sided) crib sheet.
 - Bring your ID!
- Reminder: Prof. Gittens will be taking over lectures starting on Thursday.

Answering a question from last time

• What is $(x + y)^2$?

$$x^2 + 2xy + y^2$$

- What is $(x + y)^3$? $x^3 + 3x^2y + 3xy^2 + y^3$
- What is $(x + y)^4$? $x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$

Answering a question from last time

• What is $(x + y)^2$?

$$1x^2 + 2xy + 1y^2$$

• What is $(x + y)^3$? $1x^3 + 3x^2y + 3xy^2 + 1y^3$

- What is $(x + y)^4$? $1x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + 1y^4$
- What is $(x + y)^n$?

$$\sum_{k=0}^{n} \binom{n}{k} x^k y^{n-k}$$

What we know so far

From a set of *n* objects, select *k* of them, and...

	Without replacement	With replacement
we are choosing a <u>sequence</u> .	$_{n}P_{k} = \frac{n!}{(n-k)!}$	n^k
we are choosing a <u>subset</u> .	$_{n}C_{k} = {n \choose k} = \frac{n!}{k! (n-k)!}$	$\binom{k+(n-1)}{(n-1)}$

The multinomial coefficient

- A small strand of Christmas lights has five red lights, four blue lights, and three green lights. In how many ways can they be arranged on the strand?
- Careful! This sounds like a sequence question, but consider a strand that instead had 8 white lights and 4 blue ones. How many ways could you arrange that?
 - That's just a bit string with 8 ones and 4 zeros! It would be $\binom{12}{8}$!
 - Though I advise you think of it as $\binom{12}{8}\binom{4}{4}$...

The multinomial coefficient

 A small strand of Christmas lights has five red lights, four blue lights, and three green lights. In how many ways can they be arranged on the strand?

• Start simple: How many different ways could you select the locations for the **red** lights?

After placing the red lights, how many ways could you select the locations for the blue lights?

After placing both the red and blue lights, how many ways could you select locations for the green lights?

The multinomial coefficient

- A small strand of Christmas lights has five red lights, four blue lights, and three green lights. In how many ways can they be arranged on the strand?
- Now we can apply the Product Rule: $\binom{12}{5} \cdot \binom{7}{4} \cdot \binom{3}{3}$

Quick check-in

 How many ways are there to rearrange the letters in RENSSELAER?

•
$$\binom{10}{3,2,2,1,1,1} = \frac{10!}{3! \cdot 2! \cdot 2! \cdot 1! \cdot 1!} = 151,200$$


What we know so far

From a set of *n* objects, select *k* of them, and...


	Without replacement	With replacement
we are choosing a <u>sequence</u> .	$_{n}P_{k} = \frac{n!}{(n-k)!}$	n^k
we are choosing a <u>subset</u> .	$_{n}C_{k} = \binom{n}{k} = \frac{n!}{k! (n-k)!}$	$\binom{k + (n-1)}{(n-1)}$
we are choosing a <u>sequence</u> , but some of the objects are <u>identical</u> .	$\binom{k_1+\cdots+k_r}{k_1,\ldots,k_r} = \frac{k!}{k_1!\ldots k_r!}$	

The Sum Rule

- Super basic / obvious: If a set S can be partitioned into disjoint subsets $A_1, A_2, ..., A_n$ (such that $\forall i, j \ A_i \cap A_j = \emptyset$ and $A_1 \cup A_2 \cup \cdots \cup A_n = S$), then $|S| = |A_1| + ... + |A_n|$
- What happens if the sets <u>aren't</u> disjoint? Let's start with just two sets...



Inclusion-exclusion



• Clearly we can't just say $|A \cup B| = |A| + |B|$ anymore.







- What's the problem?
 - We've double-counted the purple section we need to correct for that.

• Our rule is $|A \cup B| = |A| + |B| - |A \cap B|$


Quick check-in

- How many integers from 1 to 100 inclusive are divisible by 2 or by 5?
 - A = integers 1-100 divisible by 2
 - B = integers 1-100 divisible by 5
 - We want $|A \cup B|$.
 - $A \cap B = \text{integers } 1-100 \text{ divisible by } 2 \text{ and by } 5$
 - $|A| = 50, |B| = 20, |A \cap B| = 10$ (Why?)
 - Thus, $|A \cup B| = 50 + 20 10 = 60$

More inclusion-exclusion


What if there are three sets?

Or four sets?



Count the regions, then count how many times each region is covered by one of the main sets.

More inclusion-exclusion



- The outer sections are covered <u>once</u>, so we must start with |A| + |B| + |C|.
- The sections that are covered <u>twice</u> must be removed, giving $|A| + |B| + |C| |A \cap B| |A \cap C| |B \cap C|$.
- But this adds the center section three times, and then removes it three times, and so we must add it back in: $|A| + |B| + |C| |A \cap B| |A \cap C| |B \cap C| + |A \cap B \cap C|$
- The pattern continues if there are more sets, alternating addition and subtraction.

Less-quick check in

- How many 8-bit strings have four consecutive 0s <u>or</u> four consecutive 1s? (Note: The counting <u>before</u> you get to inclusion-exclusion will take some work.)
 - Where do the consecutive 0s start?
 - 0000xxxx: $2^4 = 16$ strings
 - $10000xxx: 2^3 = 8 \text{ strings}$
 - $x10000xx: 2^3=8 \text{ strings}$
 - $xx10000x: 2^3 = 8 \text{ strings}$
 - $xxx10000: 2^3=8 \text{ strings}$
 - So there are 48 eight-bit strings with 4+ consecutive zeros.

Less-quick check in

- How many 8-bit strings have four consecutive 0s <u>or</u> four consecutive 1s? (Note: The counting <u>before</u> you get to inclusion-exclusion will take some work.)
 - There are 48 eight-bit strings with 4+ consecutive 0s.
 - By symmetry, there are 48 eight-bit strings with 4+ consecutive 1s.
 - How many strings have <u>both</u>?
 - Just two: 00001111 and 11110000
 - That makes our final answer 48 + 48 2 = 94 strings.

The most obvious theorem in FOCS

- The Pigeonhole Principle
 - If you have more objects than places to put them, then at least one of those places will end up with two or more objects.
 - That's it. That's the whole thing.
 - It is (almost) a straightforward contraposition of the Bijection Principle.
 - If you can put two sets into 1:1 correspondence, then they are the same size.
 - If two sets are not the same size, you cannot put them into 1:1 correspondence.

 How many people do you need in a room before two of them were born in the same month?

 Every graph has a pair of vertices with the same degree.

 Any list of one hundred 27-digit positive integers will have two subsets that add up to the same value.

- How many people do you need in a room to guarantee two of them were born in the same month?
 - 13. Once you have one per month, the next person must duplicate SOMEBODY.

- Every graph with at least two vertices has a pair of vertices with the same degree.
 - With *n* vertices:
 - If the graph is connected, then the available degrees are $1 \dots n-1$.
 - If the graph is disconnected, then the available degrees are $0 \dots n-2$.

- Any list of one hundred 27-digit positive integers will have two subsets that add up to the same value.
 - The maximum value of a 27-digit number is $< 10^{28}$.
 - Therefore, the maximum sum is $< 100 \cdot 10^{28}$, or 10^{30} .
 - There are 2^{100} possible subsets.
 - $2^{100} \approx 1.26 \times 10^{30}$

Questions?

REMINDER: Exam 2 on W 2/26

Last names A-K: DCC 318

Last names L-Z: DCC 324

Bring your ID card!

