

CSCI 6971/4971 Spring 2018 Self-Assessment

Linear Algebra

Concepts: SVD, EVD, QR decompositions; Rank, Nullity; Positivity; Orthonormality; Vector norms; Frobenius norm; Spectral norm

1. What is the tightest upper bound on $|\mathbf{x}^T \mathbf{y}|$ in terms of the Euclidean norms of \mathbf{x} and \mathbf{y} ?
2. Let matrices \mathbf{A} and \mathbf{B} have the same dimensions; show that the trace of the matrix $\mathbf{A}\mathbf{B}^T$ is the same as the inner product of the vectors \mathbf{a} and \mathbf{b} obtained by stacking the columns of \mathbf{A} and \mathbf{B} .
3. Why is it that $\mathbf{A}^T \mathbf{A} + \mathbf{I}$ is invertible for any matrix \mathbf{A} ?
4. Express $\|\mathbf{A}^{-1}\|_2$ in terms of the singular values of \mathbf{A} .
5. If $\mathbf{A} = \mathbf{Q}\mathbf{R}$ is a QR decomposition, give an expression for $\mathbf{A}^T \mathbf{A}$ in terms of \mathbf{R} .
6. If $\mathbf{A} = \mathbf{U}\mathbf{S}\mathbf{V}^T$ is the full SVD of \mathbf{A} , then what are the full SVDs of $\mathbf{A}^T \mathbf{A}$ and $\mathbf{A}\mathbf{A}^T$?

7. If $\mathbf{A} = \mathbf{U}\mathbf{S}\mathbf{V}^T$ is the full SVD of \mathbf{A} , how can you read off the rank and nullity of \mathbf{A} from just \mathbf{S} ?
8. If $\mathbf{x}^T \mathbf{A}^T \mathbf{A} \mathbf{x} = 0$, what can we say about \mathbf{x} ?
9. If \mathbf{A} is symmetric (not necessarily positive-definite), how are its eigenvalue decomposition and singular value decomposition related?
10. If \mathbf{U} is a matrix with orthonormal columns, argue that $\|\mathbf{U}\mathbf{x}\|_2^2 = \|\mathbf{x}\|_2^2$.
11. Express the Frobenius norm of \mathbf{A} in terms of the trace of the matrix $\mathbf{A}^T \mathbf{A}$ and argue that it is smaller than the spectral norm of \mathbf{A} .
12. If \mathbf{A} and \mathbf{B} are matrices whose columns are respectively $\{\mathbf{a}_i\}_i$ and $\{\mathbf{b}_i\}_i$, show that $\mathbf{A}\mathbf{B}^T = \sum_i \mathbf{a}_i \mathbf{b}_i^T$.
13. Express $\|\mathbf{a}\mathbf{a}^T\|_F^2$ in terms of the Euclidean length of \mathbf{a} .

Probability

Concepts: Independence; Variance; Expectation; Total Probability; Gaussians

1. Let $p(x, y)$ be the joint pdf for two random variables X, Y in \mathbb{R}^2 ; give expressions for $p_1(x)$ and $p_2(y)$, the marginals of X and Y .
2. If X and Y are independent random variables, how is p related to p_1 and p_2 ?
3. If X and Y are independent, give an expression for the expectation of $f(x)g(y)$, where f and g are arbitrary functions (this expression should not be true in general if X and Y are not independent).
4. What can be said about $\mathbb{P}\{\mathcal{E}_1 \cap \mathcal{E}_2\}$ if the events \mathcal{E}_1 and \mathcal{E}_2 are independent?
5. If X, Y, Z are independent, what can we say about $\mathbb{E}(X+Y+Z)$? If they are not independent?
6. Let $[[\cdot]]$ be the indicator function that returns 1 if the argument is true and 0 otherwise; give a simple expression for $\mathbb{E}([[X \in \mathcal{A}]])$.
7. Use the law of total probability to argue that $\mathbb{P}\{\mathcal{E}_1\} \leq \mathbb{P}\{\mathcal{E}_1 \cap \mathcal{E}_2\} + \mathbb{P}\{\mathcal{E}_2^c\}$.

8. Give a sufficient condition for when the variance of $X + Y$ equals the sum of the variances of X and Y . Provide a proof of your claim.
9. Let \mathbf{x} be a standard multivariate Gaussian in \mathbb{R}^n , \mathbf{O} be an orthonormal matrix, and \mathcal{A} be a subset of \mathbb{R}^n . Argue that $\mathbb{P}\{\mathbf{O}\mathbf{x} \in \mathcal{A}\} = \mathbb{P}\{\mathbf{x} \in \mathbf{O}^T\mathcal{A}\}$, and evaluate the expression for the latter probability to argue that $\mathbf{O}\mathbf{x}$ is also a standard multivariate Gaussian.
10. If X_1, \dots, X_n are independent identically distributed samples drawn from a distribution with finite variance, what can we say about the distribution of their average $\frac{1}{n} \sum_{i=1}^n X_i$?