

1. Find all $z \geq z$ satisfies the LCP(A,b), where

$$A = \begin{bmatrix} 1 & 3 \\ -2 & 0 \end{bmatrix} \quad \text{and} \quad b = \begin{bmatrix} 6 \\ 4 \end{bmatrix}$$

2. Repeat the problem with $b = \begin{bmatrix} -6 \\ -4 \end{bmatrix}$

3. Repeat the problem with $b = \begin{bmatrix} -6 \\ 4 \end{bmatrix}$

$$\begin{bmatrix} -6 \\ -4 \end{bmatrix}$$

$$\begin{bmatrix} 0 \\ 0 \end{bmatrix} \perp \begin{bmatrix} -6 \\ -4 \end{bmatrix} \geq 0 \quad \text{FAIL!}$$

$$\begin{bmatrix} 0 \\ 0 \end{bmatrix} \perp \begin{bmatrix} -4 \\ -4 \end{bmatrix} = 0 \quad \text{FAIL!}$$

Try setting the r.h.s. to zero

$$Az + b = 0 \Rightarrow z = -A^{-1}b = \begin{bmatrix} 2 \\ -8/3 \end{bmatrix} \quad \text{FAIL!}$$

Try setting one r.h.s. ineq. to zero, i.e. $0 \leq \begin{bmatrix} + \\ 0 \end{bmatrix} \perp \begin{bmatrix} 0 \\ + \end{bmatrix} \geq 0$

Enforce the equations:

$$z_2 = 0, \quad z_1 + 3z_2 - 6 \geq 0 \Rightarrow z_1 \geq 6$$

Test the inequalities

$$z_1 \geq 0 \quad \text{Yes.} \quad -2z_1 - 4 \geq 0 \quad \text{No!}$$

Last chance for solution $0 \leq \begin{bmatrix} 0 \\ + \end{bmatrix} \perp \begin{bmatrix} + \\ 0 \end{bmatrix} \geq 0$

Eqs:

$$z_1 = 0, \quad -2z_1 - 4 = 0 \quad \leftarrow \text{Notice 2 equations gave us only one piece of info!}$$

Ineqs:

$$-4 \geq 0 \quad \text{Fail!}$$

We could have seen this at the start.

$$0 \leq z_1 \perp \text{~~~~~} \geq 0$$

$$0 \leq \text{~} \perp \text{~~~~~} -2z_1 - 4 \geq 0$$

Will always violate

$$0 \leq z_1 \perp -2z_1 - 4 \geq 0$$

Will always violate nonnegativity

3. Repeat the problem with $b = \begin{bmatrix} -6 \\ 4 \end{bmatrix}$.

$$0 \leq z_1 \perp z_1 + 3z_2 - 6 \geq 0$$

$$0 \leq z_2 \perp -2z_1 + 4 \geq 0$$

Notice: $z_1 = z_2 = 0$ can't work because of the -6 .

If we let z_1 & z_2 grow enough, the -6 will be balanced.

If we let z_1 grow too much we'll cause infeasibility of the second r.h. inequality.

Case 1: $\begin{bmatrix} 0 \\ 0 \end{bmatrix} \perp \begin{bmatrix} + \\ + \end{bmatrix}$ FAIL!

Case 2: $\begin{bmatrix} + \\ + \end{bmatrix} \perp \begin{bmatrix} 0 \\ 0 \end{bmatrix}$ $z = -A^{-1} \begin{bmatrix} -6 \\ 4 \end{bmatrix} = \begin{bmatrix} 2 \\ 4/3 \end{bmatrix}$ Success!

Case 3: $\begin{bmatrix} + \\ 0 \end{bmatrix} \perp \begin{bmatrix} 0 \\ + \end{bmatrix}$

$$\text{Eqs.: } z_2 = 0, z_1 + 3z_2 - 6 = 0 \Rightarrow z_1 = 6$$

$$\text{Ineqs.: } z_1 \geq 0, -2z_1 + 4 \geq 0 \text{ FAIL!}$$

Case 4: $\begin{bmatrix} 0 \\ + \end{bmatrix} \perp \begin{bmatrix} + \\ 0 \end{bmatrix}$

$$\text{Eqs. } z_1 = 0, -2z_1 + 4 = 0 \Rightarrow z_1 = 2 \text{ Contradiction!}$$

Graphical interpretation of solution


$$0 \leq \begin{bmatrix} z_1 \\ z_2 \end{bmatrix} \perp \begin{bmatrix} z_1 + 3z_2 - 6 \\ -2z_1 + 4 \end{bmatrix} \geq 0$$

$$z_1 + 3z_2 - 6 \geq 0$$

$$z_2 \geq -\frac{1}{3}z_1 + 2$$

$$-2z_1 + 4 \geq 0$$

$$z_1 \leq 2$$



$\text{SOL}(\text{LCP}(A, b))$
is unique