





The Rendering Equation & Radiosity II



An early application of radiative heat transfer in stables.

Last Time?


- Local Illumination
 - BRDF
 - Ideal Diffuse Reflectance
 - Ideal Specular Reflectance
 - The Phong Model
- Radiosity Equation/Matrix
- Calculating the Form Factors



Today

- **The Rendering Equation**
- Radiosity Equation/Matrix
- Advanced Radiosity
 - Progressive Radiosity
 - Adaptive Subdivision
 - Discontinuity Meshing
 - Hierarchical Radiosity


The Rendering Equation



$$L(x', \omega') = E(x', \omega') + \int \rho_r(\omega, \omega') L(x, \omega) G(x, x') V(x, x') dA$$

$L(x', \omega')$ is the radiance from a point on a surface in a given direction ω'


The Rendering Equation



$$L(x', \omega') = E(x', \omega') + \int \rho_r(\omega, \omega') L(x, \omega) G(x, x') V(x, x') dA$$

$E(x', \omega')$ is the emitted radiance from a point: E is non-zero only if x' is emissive (a light source)


The Rendering Equation



$$L(x', \omega') = E(x', \omega') + \int \rho_r(\omega, \omega') L(x, \omega) G(x, x') V(x, x') dA$$

Sum the contribution from all of the other surfaces in the scene


The Rendering Equation



$$L(x', \omega') = E(x', \omega') + \int \rho_x(\omega, \omega') L(x, \omega) G(x, x') V(x, x') dA$$


For each x, compute $L(x, \omega)$, the radiance at point x in the direction ω (from x to x')

The Rendering Equation




$$L(x', \omega') = E(x', \omega') + \int \rho_x(\omega, \omega') L(x, \omega) G(x, x') V(x, x') dA$$

scale the contribution by $\rho_x(\omega, \omega')$, the reflectivity (BRDF) of the surface at x'




The Rendering Equation



$$L(x', \omega') = E(x', \omega') + \int \rho_x(\omega, \omega') L(x, \omega) G(x, x') V(x, x') dA$$

For each x, compute $V(x, x')$, the visibility between x and x': 1 when the surfaces are unobstructed along the direction ω , 0 otherwise

The Rendering Equation




$$L(x', \omega') = E(x', \omega') + \int \rho_x(\omega, \omega') L(x, \omega) G(x, x') V(x, x') dA$$

For each x, compute $G(x, x')$, which describes the on the geometric relationship between the two surfaces at x and x'

Intuition about $G(x, x')$?

- Which arrangement of two surfaces will yield the greatest transfer of light energy? Why?



Questions?



Lightscape <http://www.lightscape.com>

Today


- The Rendering Equation
- Radiosity Equation/Matrix
- Advanced Radiosity
 - Progressive Radiosity
 - Adaptive Subdivision
 - Discontinuity Meshing
 - Hierarchical Radiosity

Radiosity Equation

$$L(x', \omega') = E(x', \omega') + \int \rho_{x'}(\omega, \omega') L(x, \omega) G(x, x') V(x, x') dA$$

↓ Radiosity assumption:
perfectly diffuse surfaces (not directional)

$$B_{x'} = E_{x'} + \rho_{x'} \int B_x G(x, x') V(x, x')$$



Solving the Radiosity Matrix

The radiosity of a single patch i is updated for each iteration by *gathering* radiosities from all other patches:

$$\begin{bmatrix} B_1 \\ B_2 \\ \vdots \\ B_i \\ \vdots \\ B_n \end{bmatrix} = \begin{bmatrix} E_1 \\ E_2 \\ \vdots \\ E_i \\ \vdots \\ E_n \end{bmatrix} + \rho_i F_{i1} \rho_1 F_{12} \dots \rho_n F_{ni} \begin{bmatrix} B_1 \\ B_2 \\ \vdots \\ B_i \\ \vdots \\ B_n \end{bmatrix}$$


↑ Radiosity values on iteration $t+1$ ↑ Radiosity values on iteration t

This method is fundamentally a Gauss-Seidel relaxation

Today


- The Rendering Equation
- Radiosity Equation/Matrix
- Advanced Radiosity
 - Progressive Radiosity
 - Adaptive Subdivision
 - Discontinuity Meshing
 - Hierarchical Radiosity

Stages in a Radiosity Solution




Progressive Refinement

- Goal: Provide frequent and timely updates to the user during computation
- Key Idea: Update the entire image at every iteration, rather than a single patch
- How? Instead of summing the light received by one patch, distribute the radiance of the patch with the most *undistributed radiance*.




Reordering the Solution for PR

Shooting: the radiosity of all patches is updated for each iteration:


$$\begin{bmatrix} B_1 \\ B_2 \\ \vdots \\ B_n \end{bmatrix} = \begin{bmatrix} B_1 \\ B_2 \\ \vdots \\ B_n \end{bmatrix} + \begin{bmatrix} \rho_1 F_{1i} & \dots \\ \rho_2 F_{2i} & \dots \\ \vdots & \vdots \\ \rho_n F_{ni} & \dots \end{bmatrix} \begin{bmatrix} \vdots \\ \vdots \\ \vdots \\ B_i \end{bmatrix}$$


This method is fundamentally a Southwell relaxation

Progressive Refinement w/out Ambient Term



Progressive Refinement with Ambient Term



Questions?




Lightscape <http://www.lightscape.com>

Today

- The Rendering Equation
- Radiosity Equation/Matrix
- Advanced Radiosity
 - Progressive Radiosity
 - Adaptive Subdivision
 - Discontinuity Meshing
 - Hierarchical Radiosity


Increasing the Accuracy of the Solution

What's wrong with this picture?



- Image quality is a function of patch size
- Compute a solution on a uniform initial mesh, then refine the mesh in areas that exceed some error tolerance:
 - shadow boundaries
 - other areas with a high radiosity gradient

Adaptive Subdivision of Patches



Coarse patch solution
(145 patches)


Improved solution
(1021 subpatches)

Adaptive subdivision
(1306 subpatches)


Discontinuity Meshing

- Limits of umbra and penumbra

- Captures nice shadow boundaries
- Complex geometric computation to construct mesh




Discontinuity Meshing



"Fast and Accurate Hierarchical Radiosity Using Global Visibility"
Durand, Drettakis, & Puech 1999


Hierarchical Radiosity

- Group elements when the light exchange is not important
 - Breaks the quadratic complexity
 - Control non trivial, memory cost



Practical Problems with Radiosity

- Meshing
 - memory
 - robustness
- Form factors
 - computation
- Diffuse limitation
 - extension to specular takes too much memory



Cow-cow form factor?

Questions?



Lightscape <http://www.lightscape.com>

Reading for Today:

- “A Two-Pass Solution to the Rendering Equation: A Synthesis of Ray Tracing and Radiosity Methods”
Wallace, Cohen, & Greenberg, SIGGRAPH 1987



direct illumination
(standard raytracing)


indirect illumination
(standard radiosity)

full solution

- *Optional Reading:* “The Rendering Equation”
Kajiya, SIGGRAPH 1986

Reading for Friday 3/4:

- “Rendering Fake Soft Shadows with Smoothies”, Chan & Durand, 2003.



shadow volumes

shadow volumes w/ “smoothies”

Post a comment or question on the LMS
discussion by 10am on Friday 3/4