

Foundations of Computer Science

Lecture 18

Random Variables

Measurable Outcomes

Probability Distribution Function

Bernoulli, Uniform, Binomial and Exponential Random Variables



① Independence.

- ▶ Using independence to estimate complex probabilities.

② Coincidence.

- ▶ FOCS-twins.
- ▶ The birthday paradox.
- ▶ Application to hashing.

③ Random walks and gambler's ruin.

Today: Random Variables

- 1 What is a random variable?
- 2 Probability distribution function (PDF) and Cumulative distribution function (CDF).
- 3 Joint probability distribution and independent random variables
- 4 Important random variables
 - Bernoulli: indicator random variables.
 - Uniform: simple and powerful. An equalizing force.
 - Binomial: sum of independent indicator random variables.
 - Exponential: the waiting time to the first success.

A Random Variable is a “Measurable Property”

Temperature: “measurable property” of random positions and velocities of molecules.

Toss 3 coins.

number-of-heads(HTT) = 1;
all-tosses-match(HTT) = 0.

	SAMPLE SPACE Ω								
ω	HHH	HHT	HTH	HTT	THH	THT	TTH	TTT	
$P(\omega)$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{8}$	
$\mathbf{X}(\omega)$	3	2	2	1	2	1	1	0	← number of heads
$\mathbf{Y}(\omega)$	1	0	0	0	0	0	0	1	← matching tosses
$\mathbf{Z}(\omega)$	8	2	2	$\frac{1}{2}$	2	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{8}$	← H: double your money T: halve your money

Can use random variables to define events:

$$\begin{aligned}
 \{\mathbf{X} = 2\} &= \{\text{HHT, HTH, THH}\} & \mathbb{P}[\mathbf{X} = 2] &= \frac{3}{8} \\
 \{\mathbf{X} \geq 2\} &= \{\text{HHH, HHT, HTH, THH}\} & \mathbb{P}[\mathbf{X} \geq 2] &= \frac{1}{2} \\
 \{\mathbf{Y} = 1\} &= \{\text{HHH, TTT}\} & \mathbb{P}[\mathbf{Y} = 1] &= \frac{1}{4} \\
 \{\mathbf{X} \geq 2 \text{ AND } \mathbf{Y} = 1\} &= \{\text{HHH}\} & \mathbb{P}[\mathbf{X} \geq 2 \text{ AND } \mathbf{Y} = 1] &= \frac{1}{8}
 \end{aligned}$$

Probability Distribution Function (PDF)


$$\underbrace{\{\text{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT}\}}_{\Omega} \xrightarrow{\mathbf{X}} \underbrace{\{3, 2, 1, 0\}}_{\mathbf{X}(\Omega)}$$

Each *possible* value x of the random variable \mathbf{X} corresponds to an event,

x	0	1	2	3
Event	{TTT}	{HTT, THT, TTH}	{HHT, HTH, THH}	{HHH}

For each $x \in \mathbf{X}(\Omega)$, compute $\mathbb{P}[\mathbf{X} = x]$ by adding the outcome-probabilities,

x	possible values $x \in \mathbf{X}(\Omega)$			
	0	1	2	3
$P_{\mathbf{X}}(x)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$















Probability Distribution Function (PDF). The probability distribution function $P_{\mathbf{X}}(x)$ is the probability for the random variable \mathbf{X} to take value x ,

$$P_{\mathbf{X}}(x) = \mathbb{P}[\mathbf{X} = x].$$

PDF for the Sum of Two Dice

Probability Space


Die 2 Value		$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
		$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
		$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
		$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
		$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
		$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
							
		Die 1 Value					

$\mathbf{X} = 9$ has four outcomes,

$$\mathbb{P}[\mathbf{X} = 9] = 4 \times \frac{1}{36} = \frac{1}{9}.$$

Possible sums are $\mathbf{X} \in \{2, 3, \dots, 12\}$ and the PDF is

x	2	3	4	5	6	7	8	9	10	11	12
$P_{\mathbf{X}}(x)$	$\frac{1}{36}$	$\frac{1}{18}$	$\frac{1}{12}$	$\frac{1}{9}$	$\frac{5}{36}$	$\frac{1}{6}$	$\frac{5}{36}$	$\frac{1}{9}$	$\frac{1}{12}$	$\frac{1}{18}$	$\frac{1}{36}$



Joint PDF: More Than One Random Variable

ω	SAMPLE SPACE Ω								
	HHH	HHT	HTH	HTT	THH	THT	TTH	TTT	
$P(\omega)$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{8}$	
$\mathbf{X}(\omega)$	3	2	2	1	2	1	1	0	← number of heads
$\mathbf{Y}(\omega)$	1	0	0	0	0	0	0	1	← matching tosses

$$\begin{aligned} \mathbb{P}[\mathbf{X} = 0, \mathbf{Y} = 0] &= 0 \\ \mathbb{P}[\mathbf{X} = 1, \mathbf{Y} = 0] &= \frac{3}{8}. \end{aligned}$$

$$P_{\mathbf{XY}}(x, y) = \mathbb{P}[\mathbf{X} = x, \mathbf{Y} = y].$$

$$\mathbb{P}[\mathbf{X} + \mathbf{Y} \leq 2] = 0 + \frac{3}{8} + \frac{3}{8} + \frac{1}{8} + 0 = \frac{7}{8}.$$

$$\mathbb{P}[\mathbf{Y} = 1 \text{ AND } \mathbf{X} + \mathbf{Y} \leq 2] = \frac{1}{8} + 0 = \frac{1}{8}.$$

$$\begin{aligned} \mathbb{P}[\mathbf{Y} = 1 \mid \mathbf{X} + \mathbf{Y} \leq 2] &= \frac{\mathbb{P}[\mathbf{Y}=1 \text{ AND } \mathbf{X}+\mathbf{Y} \leq 2]}{\mathbb{P}[\mathbf{X}+\mathbf{Y} \leq 2]} \\ &= \frac{1/8}{7/8} = \frac{1}{7} \end{aligned}$$

		\mathbf{X}				row sums
		0	1	2	3	
\mathbf{Y}	0	0	$\frac{3}{8}$	$\frac{3}{8}$	0	$\frac{3}{4}$
	1	$\frac{1}{8}$	0	0	$\frac{1}{8}$	$\frac{1}{4}$
column sums		$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$	

$$P_{\mathbf{Y}}(y) = \sum_{x \in \mathbf{X}(\Omega)} P_{\mathbf{XY}}(x, y)$$

$$P_{\mathbf{X}}(x) = \sum_{y \in \mathbf{Y}(\Omega)} P_{\mathbf{XY}}(x, y)$$

Independent Random Variables

Independent Random Variables measure unrelated quantities. The joint-PDF is *always* the product of the marginals.

$$P_{\mathbf{XY}}(x, y) = P_{\mathbf{X}}(x)P_{\mathbf{Y}}(y) \quad \text{for all } (x, y) \in \mathbf{X}(\Omega) \times \mathbf{Y}(\Omega).$$

Our \mathbf{X} and \mathbf{Y} are *not* independent,


		\mathbf{X}				
		0	1	2	3	
\mathbf{Y}	0	0	$\frac{3}{8}$	$\frac{3}{8}$	0	$\frac{3}{4}$
	1	$\frac{1}{8}$	0	0	$\frac{1}{8}$	$\frac{1}{4}$
		$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$	

		\mathbf{X}				
		0	1	2	3	
\mathbf{Y}	0	$\frac{3}{32}$	$\frac{9}{32}$	$\frac{9}{32}$	$\frac{9}{32}$	$\frac{3}{4}$
	1	$\frac{1}{32}$	$\frac{3}{32}$	$\frac{3}{32}$	$\frac{1}{32}$	$\frac{1}{4}$
		$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$	

Practice: Exercise 18.4, Pop Quizzes 18.5, 18.6.

Cumulative Distribution Function (CDF)

x	0	1	2	3
$P_{\mathbf{X}}(x)$	$\frac{1}{8}$	$\frac{3}{8}$	$\frac{3}{8}$	$\frac{1}{8}$
$\mathbb{P}[\mathbf{X} \leq x]$	$\frac{1}{8}$	$\frac{4}{8}$	$\frac{7}{8}$	$\frac{8}{8}$



Cumulative Distribution Function (CDF). The cumulative distribution function $F_{\mathbf{X}}(x)$ is the probability for the random variable \mathbf{X} to be at most x ,

$$F_{\mathbf{X}}(x) = \mathbb{P}[\mathbf{X} \leq x].$$

Bernoulli Random Variable: Binary Measurable $(0, 1)$

Two outcomes: coin toss, drunk steps left or right, etc. \mathbf{X} *indicates* which outcome,

$$\mathbf{X} = \begin{cases} 1 & \text{with probability } p; \\ 0 & \text{with probability } 1 - p. \end{cases}$$

Can add Bernoullis. Toss n independent coins. \mathbf{X} is the number of H.

$$\mathbf{X} = \mathbf{X}_1 + \mathbf{X}_2 + \cdots + \mathbf{X}_n.$$

\mathbf{X} is a sum of Bernoullis, each \mathbf{X}_i is an independent Bernoulli.

Drunk makes n steps. Let \mathbf{R} be the number of right steps

$$\mathbf{R} = \mathbf{X}_1 + \mathbf{X}_2 + \cdots + \mathbf{X}_n.$$

\mathbf{R} is a sum of Bernoullis. $\mathbf{L} = n - \mathbf{R}$ and the final position \mathbf{X} is:


$$\mathbf{X} = \mathbf{R} - \mathbf{L} = 2\mathbf{R} - n = 2(\mathbf{X}_1 + \mathbf{X}_2 + \cdots + \mathbf{X}_n) - n.$$

Uniform Random Variable: Every Value Equally Likely

n possible values $\{1, 2, \dots, n\}$, each with probability $\frac{1}{n}$:

$$P_X(k) = \frac{1}{n}, \quad \text{for } k = 1, \dots, n.$$

Roll of a 6-sided *fair* die $\sim \mathbf{U}[6]$. (Uniform on $\{1, \dots, 6\}$)



Example: Matching game (uniform is an equalizer in games of strategy).
GR will pick a path to relieve you of your lunch money.
If you pick your path uniformly, you win half the time.



Example 18.2: Guessing Larger or Smaller

I pick two numbers from $\{1, \dots, 5\}$, as I please. I *randomly* show you one of the two, x .
You must guess if x is the larger or smaller of my two numbers.

You always say smaller: you win $\frac{1}{2}$ the time.

You say smaller if $x \leq 3$ and larger if $x > 3$. I pick numbers 1,2: you win $\frac{1}{2}$ the time.

You have a strategy which wins *more* than $\frac{1}{2}$ the time, and I *cannot* prevent it!

Binomial Random Variable: Sum of Bernoullis

\mathbf{X} = number of heads in n independent coin tosses with probability p of heads,

$$\mathbf{X} = \mathbf{X}_1 + \dots + \mathbf{X}_n.$$

← sum of n independent Bernoullis, $\mathbf{X}_i \sim \text{Bernoulli}(p)$

$n=5, \mathbf{X}=3$:

HHHTT HHTTH HTTHH TTHHH HHTHT
HTHTH THTHH HTHHT THTHT THHHT

← each has probability $p^3(1-p)^2$ (independence)


$$\mathbb{P}[\mathbf{X} = 3 \mid n = 5] = 10p^3(1-p)^2$$

← add outcome probabilities

In general, $\binom{n}{k}$ sequences with k heads.

Each has probability $p^k(1-p)^{n-k}$, so

$$\mathbb{P}[\mathbf{X} = k \mid n] = \binom{n}{k} p^k(1-p)^{n-k}.$$



Binomial Distribution. \mathbf{X} is the number of successes in n independent trials with success probability p on each trial: $\mathbf{X} = \mathbf{X}_1 + \dots + \mathbf{X}_n$ where $\mathbf{X}_i \sim \text{Bernoulli}(p)$.

$$P_{\mathbf{X}}(k) = B(k; n, p) = \binom{n}{k} p^k(1-p)^{n-k}.$$


Example: guessing correctly on the multiple choice quiz: $n = 15$ questions, 5 choices ($p = \frac{1}{5}$).

number correct, k	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
probability	0.035	0.132	0.231	0.250	0.188	0.103	0.043	0.014	0.003	7×10^{-4}	10^{-4}	10^{-5}	10^{-6}	~ 0	~ 0	~ 0

chances of passing are $\approx 0.4\%$


Exponential Random Variable: Waiting Time to Success

Let p be the probability to succeed on a trial.



$$\mathbb{P}[t \text{ trials}] = \mathbb{P}[F^{t-1}S] = (1 - p)^{t-1}p$$

$$P_X(t) = (1 - p)^{t-1}p = \underbrace{\frac{p}{1 - p}}_{\beta} \times (1 - p)^t = \beta(1 - p)^t.$$



Example: 3 people randomly access the wireless channel. Success only if exactly one is attempting.

Try every timestep \rightarrow no one succeeds. Everyone tries $\frac{1}{3}$ the time (randomly).
 Success probability for *someone* is $\frac{4}{9}$. Success probability for *you* is $\frac{4}{27}$.

wait, t	1	2	3	4	5	6	7	8	9	10	11	...
$\mathbb{P}[\text{someone succeeds}]$	0.444	0.247	0.137	0.076	0.042	0.024	0.013	0.007	0.004	0.002	0.001	...
$\mathbb{P}[\text{you succeed}]$	0.148	0.126	0.108	0.092	0.078	0.066	0.057	0.048	0.051	0.035	0.030	...